

SPECTROSCOPIC CHARACTERIZATION OF HUMIC SUBSTANCES OF ANTHROPOGENIC SOILS DERIVED FROM TERRA ROSSA

Aleksandra BENSA^{1*}, Boško MILOŠ², Nikolina JURKOVI BALOG¹, Ivana TORI¹

¹University of Zagreb Faculty of Agriculture, Svetošimunska 25, 10 000 Zagreb, Croatia

²Institute for Adriatic Crops and Karst Reclamation, Put Duilova 11, 21 000 Split, Croatia

*Corresponding author: abensa@agr.hr

ABSTRACT

The aim of the study was spectroscopic characterization of humic substances and evaluation of humus quality of anthropogenic soils derived from Terra Rossa. The study was conducted on 15 soil samples collected from top-soil horizon of olive groves in Middle Dalmatia (Croatia). Total organic carbon was determined according to Walkley-Black method (1934) and humic substances isolation following procedure given by Schnitzer (1982). Spectroscopic characterization of isolated humic substances was carried out by measuring absorbance in VIS spectral range 400-700 nm and optical indices ($Q_{4/6}$, $E_{4/6}$) were calculated. SOC content in top-soil samples varied from 1.45% to 4.21% with mean value of 2.98%. The absorption spectrum showed monotonous decrease of absorbance from 400 to 700 nm for all samples. The optical indices $E_{4/6}$ and $Q_{4/6}$ varied from 3.58 to 5.05 and from 3.91 to 5.04, respectively and indicated differences of humus quality. The optical index below 4, which implies a high quality of humus, was determined in six samples. The low humus quality (optical index above 4) was determined in other nine samples. The SOC content has significant positive correlation with optical indices $Q_{4/6}$ and $E_{4/6}$. Soils with higher organic carbon content have lower humus quality.

Key words: *VIS spectrum, $E_{4/6}$, humus quality, Dalmatia.*

INTRODUCTION

Humus defines key soil characteristics, its fertility and it is an indicator of the processes occurring in soil. Therefore, understanding of its content and quality is important for the sustainable land management, especially in Mediterranean karst region. Humic substances (HS) constitute a major fraction (60–70%) of soil organic matter and those are possibly the most abundant of naturally occurring organic macromolecules on the Earth ($2-3 \times 10^{10}$ t), (Jones and Bryan, 1998). Humic substances differ in molecular weight, elemental composition, acidity and cation exchange capacity and can be classified into three major groups according to their solubility; humic acid, fulvic acid and humins. The humic acids fraction consists of hydroxyphenols, hydroxybenzoic acids and other aromatic structures

with linked peptides, amino compounds and fatty acids. Fulvic acids are typically composed of a variety of phenolic and benzene carboxylic acids that are held together by hydrogen bonds to form stable polymer structures. The low molecular weight of fulvic acid has higher oxygen but lower carbon content than humic acid. There are also more acidic functional groups particularly -COOH in fulvic acid molecule (Schnitzer and Khan, 1978).

Non-destructive spectroscopic methods ensure valuable informations on molecular structure, chemical and functional properties of humic substances (Chen et al, 1977). Therefore, new approaches of spectrometry that include a wide variety of techniques (UV-VIS, DRIFT, SFS, and ^{13}C -NMR) have been successfully applied to the study of HS chemical composition and structure (Pospišilova et al, 2008; Milori et al., 2002; Sierra et al., 2005). Humic substances generally show strong absorbance in the UV-VIS range (from 190 to 700 nm) because of the presence of aromatic chromophores and other organic compounds (Rupiasih and Vidyasagar, 2007). Stevenson (1982) has shown that absorption of humic substances on wave length of 465 nm is equal to absorption of light form components linked to initial phases of humification (young humic substances – fulvo acid) and absorption of light on wave length of 665 nm refers to well humified components - humic acid. Optical index $E_{4/6}$ calculated as ratio of optical absorbance at 465 nm versus 665 nm for humic substances in solution is often used for evaluation of humus quality. Generally, lower molecular weight and lower degree of condensation of aromatic structures in humic substances show higher values of $E_{4/6}$ than humic substances with a high degree of humification (Orlov, 1985).

The objective of the study was spectroscopic characterization of humic substances and evaluation of humus quality of anthropogenic soils derived from Terra Rossa.

MATERIAL AND METHODS

The study was conducted on 15 soil samples collected in three olive groves area in Middle Dalmatia (Croatia) including: the island of Bra (43°22'00" N, 16°38'27" E), Marina (43°31'40" N, 16°09'31" E) and Primošten (43°34'47" N, 15°56'38" E). Within each selected area five average soil samples from the depth 0-25cm were collected. Investigated soils are anthropogenic soils derived from Terra rossa, shallow, skeletal on limestones and dolomites (Škori et al., 1975). According to IUSS Working Group (WRB, 2014) studied soils can be identified as Chromic Leptic Skeletic Cambisol (Clayic Colluvic). Soil samples were prepared for analysis of physical and chemical properties according to HRN ISO 11464:2009. Soil pH was determined according to HRN ISO 10390:2005, soil organic carbon was determined according to Walkley-Black method (1934), available phosphorus and potassium were determined according to Egner et al. (1960) and particle size distribution was made by HRN ISO 11277:2009.

The isolation of soil humic substances (HS) was made by Schnitzer method (1982). 5 g of air dried soil sample was sieved at the mesh size of 1 mm and extracted with solution of 0.1 M NaOH + 0.1 M $\text{Na}_4\text{P}_2\text{O}_7$. The mixture was shaken mechanically for 24 h at room temperature. The supernatant solution was then separated from the

residue by centrifugation at 4000 rpm for 20 min. The alkaline extract was acidified with concentrated H_2SO_4 to pH~1, allowed to stand for 24 h at room temperature to obtain the complete precipitation of humic acid (HA). The precipitated HA was separated from fulvic acid (FA) by repeating three times the following: centrifugation at 4000 rpm for 20 min, removal of the residue, washing the HA with 0.05 M H_2SO_4 solution. Finally, the centrifuged HA were dissolved in a minimal volume of 0.1 M NaOH and brought to dryness in a drying oven at 60°C. VIS spectra were measured by Shimadzu UV 1700 spectrometer in the range of 400-700 nm. Optical indices $E_{4/6}$ and $Q_{4/6}$ were determined as the absorbance ratio $A_{465/665}$ and $A_{400/600}$, respectively (Orlov, 1985; Szajdak et al., 2006).

RESULTS AND DISCUSSION

Descriptive statistics for basic soil properties given in Table 1. shows that investigated soils are weakly acid to neutral, non-calcareous to slightly calcareous and variably supplied with physiologically active nutrients. Soil supply with P_2O_5 varies from low to moderate, showing very high coefficient of variation (CV). Investigated soils are moderate to high supplied with K_2O and showing lower coefficient of variation. Soil organic carbon content varies from low to medium, in average medium (Table 1). These data are consistent with research of Miloš and Bensa (2012) which determined similar values of SOC content in anthropogenic soils derived from Terra Rossa (0.17-3.73%) in Dalmatia. Although, only less than a quarter (24.6%) of southern European top-soils contain medium to high (>2%) amounts of SOC (Zdruli et al, 2004). Higher level of SOC content in our research, compared to averages in European soils, can be related to land management and particularities of soils formed on the limestones and dolomites in Mediterranean region (high stoniness and rockiness and variable soil depth). Investigated soils are dominantly silty clay and silty clayey loam.

Table 1. Descriptive statistics for basic soil properties

Soil property	Mean	Median	Min.	Max.	Range	Std. dev.	#CV(%)
pH H_2O	7.62	7.65	6.53	8.02	1.49	0.40	5.25
pH KCl	6.67	6.87	5.36	7.18	1.82	0.51	7.63
$CaCO_3$ (%)	3.02	2.57	0.00	7.90	7.90	2.19	72.66
P_2O_5 mg/100 g	4.37	2.50	0.05	14.13	14.08	4.49	102.71
K_2O mg/100 g	35.89	34.20	18.00	69.38	51.38	12.01	33.46
SOC %	2.98	3.24	1.46	4.23	2.77	0.96	32.3
Coarse sand (%)	22.89	23.60	15.30	28.00	12.70	3.73	16.31
Fine sand (%)	6.64	7.00	0.30	15.10	14.80	4.48	67.42
Silt (%)	27.38	29.60	13.70	37.00	23.30	6.58	24.03
Clay (%)	43.09	43.00	31.00	61.80	30.80	9.04	20.99

#CV (%), coefficient of variation

The absorption spectrums of humic substances (HS) extracted from the studied soils show monotonous decrease of absorbance in the range 400-700 nm, Figure 1. The steeper declines of curves imply domination of aromatic structures over aliphatics in humic substances (Pospišilova and Fasurova 2009; Fasurova and Pospišilova 2011; Milori et al. 2002). The steepest declines of curves were established in the VIS spectra of humic substances extracted from soils with the lowest SOC content. That means, soils with lower SOC content have more condensed aromatic structures than aliphatics in humic substances and higher humus quality.




Figure 1. VIS spectra of extracted soil humic substances (HS) from top-soil horizon of investigated soils

Therefore the ratio between the absorbance at 465 nm and 665 nm ($E_{4/6}$) and 400 and 600 nm ($Q_{4/6}$) are frequently used to characterize composition of organic matter and evaluation of its humification degree. The $E_{4/6}$ ratio decreases when the condensation and aromaticity of the humic substances rise and with increasing molecular weight (Stevenson, 1994; Fuentes et al., 2006), which is typical for more mature, more evolved organic materials, and is therefore useful as a humification indicator. The mean values of optical indices $E_{4/6}$ and $Q_{4/6}$ of humic substances isolated from investigated soils were 4.22 and 4.42 respectively, Table 2. The optical indices are characterized with short range and small coefficients of variation (CV %). These indices have lower coefficient of variation than SOC (Table 1), implying higher heterogeneity of humus quantity than quality (Table 2).

Table 2. Descriptive statistics for optical indices

Optical index	Mean	Min.	Max.	Range	St. dev.	#CV(%)
$E_{4/6}$	4.22	3.58	5.05	1.47	0.50	11.74
$Q_{4/6}$	4.42	3.91	5.04	1.13	0.33	7.41

#CV (%), coefficient of variation

Generally, optical indices values above 4 indicate presence of more aliphatic and fewer aromatic compounds and lower humus quality. From a total of 15 samples, 6 have optical index below 4, which implies a high quality of humus. The other nine samples have optical index above 4 and low humus quality. The low humus quality were determined in soil samples with SOC content above 2%. These results shows that in anthropogenic soils developed from Terra Rossa under olive groves dominated low humus quality. This probably relates with specific soil management (addition organic matter), leading to increase of SOC content and higher proportion of aliphatic compounds in humic substances (higher E_4/E_6 ratio - lower humus quality).

These results are consistent with investigation (olak and Martinovi , 1975.) which have found similar values of optical index of humic substances ($E_{4/6}$ 3.78-5.02) isolated from top-soil of Terra Rossa in Middle Dalmatia. Senesi et al (1999) also established high values of $E_{4/6}$ (4.6-4.9) in Terra rossa under olive groves in Mediterranean region.

Relations between SOC content and optical indices $Q_{4/6}$ and $E_{4/6}$ were fitted with linear type of equation: $SOC \% = -3.3522 - 0.5995 * x + 2.1249 * y$, Figure 2.




Figure 2. 3D surface plot of SOC against optical indices $Q_{4/6}$ and $E_{4/6}$

The 3D surface plot of SOC against optical indices $Q_{4/6}$ and $E_{4/6}$ (Figure 2) shows that increase of SOC content increases the value of the indices $Q_{4/6}$ and $E_{4/6}$. The strength of these relations established with linear correlation shows that SOC content and the optical indices ($Q_{4/6}$ and $E_{4/6}$) are significantly positively correlated, Table 3. The optical index $E_{4/6}$ better correlates with SOC content than $Q_{4/6}$ due to possible inaccuracies in measurements when values of absorbance are up to 2 % (around 400 nm).

Table 3. Correlations of SOC content and optical indices $E_{4/6}$ and $Q_{4/6}$

	SOC %	$Q_{4/6}$	$E_{4/6}$
SOC %	1.00	0.76*	0.92*
$Q_{4/6}$		1.00	0.88*
$E_{4/6}$			1.00

* Marked correlations are significant at $p < 0,050$

CONCLUSION

Our research showed domination of low humus quality in top-soil horizon of anthropogenic soils developed from Terra Rossa under olive groves in Middle Dalmatia, Croatia. The humus quality indicators (optical indices $Q_{4/6}$ and $E_{4/6}$) significantly correlate with SOC content. The soils with higher SOC content have lower humus quality.

REFERENCES

- Chen Y., Senesi N., Schnitzer M. (1977). Information Provided on Humic Substances by E4/E6 Ratios. SSSAJ 41 (2): 352-358.
- olak A., Martinovi J. (1975). Basic Soil Map of Croatia at 1:50 000, map sheet Šibenik 1. Project Council for the Soil Map of the Republic of Croatia, Zagreb.
- Egner H., Riehm H., Domingo W.R. (1960): Studies concerning the chemical analysis of soils as background for soil nutrient assessment. II. Chemical extracting methods to determinate the phosphorous and potassium content of soil. Kungliga Lant-u brukshögskolans annaler, 26: 199–215.
- Fasurova N., Pospíšilova L. (2011). Spectroscopic Characteristics of Humates Isolated from Different Soils. Soil & Water Res. 6 (3): 147–152.
- Fuentes, M., González-Gaitano, G., García-Mina, J.M. (2006) The usefulness of UV-visible and fluorescence spectroscopies to study the chemical nature of humic substances from soils and composts. Org. Geochem. 37, 1949-1959.
- HRN ISO 11464:2009. Soil quality - Pre-treatment of samples for physical-chemical analysis. International standard. Zagreb Croatia: Croatian Standards Institute
- HRN ISO 10390:2005. Soil quality - Determination of pH. International standard Zagreb Croatia: Croatian Standards Institute
- HRN ISO 11277:2009. Soil quality – Determination of particle size distribution (mechanical composition) in mineral soil – Method by sieving and sedimentation. International standard. Zagreb Croatia: Croatian Standards Institute
- IUSS Working Group WRB (2014). World reference base for soil resources 2014 - International soil classification system for naming soils and creating legends for soil maps. World Soil Resources Reports No. 106, FAO, Rome
- Jones M.N., Bryan N.D. (1998) Colloidal properties of humic substances. Adv. Colloid. Interface Sci. 78: 1-48.

- Milori D.M.B.P., Martin-Neto L., Bayer C., Mielniczuk J., Bagnato V.S. (2002). Humification degree of soil humic acid determined by fluorescence spectroscopy. *Soil Science*, 167: 739–749.
- Miloš B., Bensa A. (2012) Soil humus analysis by the reflectance spectroscopy, Proceedings of 47. Croatian and 7. International Symposium of Agronomy, 13.-17. February 2012., Opatija,
- Orlov D.S. (1985). *Soils Chemistry*. Moscow State University, Moscow
- Pospišilova L., Fasurova N., Baran ikova G., Liptaj T. (2008). Spectral characteristics of humic acids isolated from south Moravian lignite and soils. *Petroleum & Coal* 50/2: 30-36.
- Pospišilova L., Fasurova N. (2009). Spectroscopic Characteristics of Humic Acids Originated in Soils and Lignite. *Soil & Water Res.* 4 (4): 168–175.
- Rupiasih N.N., Vidyasagar P.B. (2007). Humic Substances: structure, function, effects and applications. *International Journal of Environment and Pollution.* 5(2):39-47.
- Schnitzer M. (1982). Organic matter characterization. In: *Methods of Soil Analysis Part 2 - Chemical and Microbiological Properties* (Weaver R.W., Angle J.S., Bottomley P.S.), Madison, USA, No. 9. 2nd Edition
- Schnitzer M., Khan S. U. (1978). *Soil Organic Matter*. Elsevier, New York
- Senesi N, Brunetti G, Loffredo E, Miano T.M. (1999) Abiotic catalytic humification of organic matter in olive oil mill wastewater, *Understanding Humic Substances – Advanced methods, Properties and Applications*, Boston, USA
- Sierra M.M.D., Giovanela M., Parlanti E., Soriano-Sierra E.J. (2005). Fluorescence fingerprint of fulvic and humic acids from varied origins as viewed by single-scan and excitation/emission matrix techniques. *Chemosphere* 58: 715-733.
- Stevenson F.J. (1982). *Humus Chemistry – Genesis, Composition, Reactions*. J. Wiley Interscience Publication, New York
- Stevenson, F.J. (1994). *Humus chemistry: genesis, composition, reactions*; Wiley, USA
- Szajdak L., Maryganova M., Tychinskaja L. (2006). Particularities of the chemical structure of humic acids from soils under shelterbelts of different age and adjoining cultivated fields. In: Frimmel F.H., Abbt-Braun G. (Eds): *Proc. 13th Meeting of the International Humic Substances Society*. Karlsruhe, 513–516.
- Škoric, A., Filipovski, G., iri , M. (1985). Classification of Yugoslav soils (in Serbo-Croatian). Department of Sciences and Arts of Bosnia and Herzegovina, Sarajevo.
- Zdruli P, Jones RJA, Montanarella L. (2004). *Organic Matter in the Soils of Southern Europe*. European Soil Bureau Technical Report, EUR 21083 EN, Luxembourg
- Walkley A, Black I.A (1934). An examination of the method for determining soil organic matter and a proposed modification of the chromic acid method, *Soil Science* 37, 29– 38.