

## Wheat Yield and Weediness under Different Rate of Nitrogen Fertilization

Desimir Knežević<sup>1</sup>, Dimitrije Marković<sup>2</sup>,  
Sretenka Marković<sup>2</sup>, Danijela Kondić<sup>2</sup>

<sup>1</sup> University of Priština, Faculty of Agriculture, Kosovska Mitrovica-Lešak, Serbia  
<sup>2</sup>Faculty of Agriculture, University of Banja Luka, Bosnia and Herzegovina

### Abstract

In this study we tested yield and weediness in wheat cultivars Lazarica and Julija Mono grown under different regime of mineral nutrition. This investigation was conducted at the long term experiment on control variant (without fertilizers), second variant with nitrogen application (two sub variants with different rate of N) 2N<sub>1</sub> (80 kg N ha<sup>-1</sup>) and 2N<sub>2</sub> (120 kg N ha<sup>-1</sup>), third variant with NP fertilizer with two different rate of N application: 3N<sub>1</sub> (80 kg N ha<sup>-1</sup> + 100kg P ha<sup>-1</sup>) and 3N<sub>2</sub> (120 kg N ha<sup>-1</sup> + 100kg P ha<sup>-1</sup>) and fourth variant 4N<sub>1</sub> (80 kg N ha<sup>-1</sup> + 60kg P ha<sup>-1</sup>) and 4N<sub>2</sub> (120 kg N ha<sup>-1</sup> + 60kg P ha<sup>-1</sup>). Weediness varies in dependence of rate and variant of fertilization. The weed species *Consolida regalis* Gray, *Convolvulus arvensis* L., *Polygonum aviculare* L., *Cirsium arvense* L., *Agropyrum repens* L., *Polygonum convolvulus* L. was dominant on fertilized plots. The analyzed variability of grain yield depends of applied fertilizer variants. The lowest grain yield was found on control variant in both wheat cultivar Lazarica (1880kg ha<sup>-1</sup>), Julija Mono (1740kg ha<sup>-1</sup>) while the highest grain yield on variant 3N<sub>2</sub> (120 kg N ha<sup>-1</sup> + 100kg P ha<sup>-1</sup>) Lazarica (3724kg ha<sup>-1</sup>) Julija Mono (4990kg ha<sup>-1</sup>) and low weediness.

**Key words:** plant competitiveness, weediness, wheat, environment, grain yield, rate of fertilizer

### Introduction

Wheat grain yield and yield components varied in dependence of crop nutrition, pest protection (Jolánkai et al., 2006) weediness and meteorological conditions (Márton, 2008). For achievement of high grain yield and quality it is necessary to optimize scientific farming measure like soil tillage (Sepp et al., 2009), optimal time and density of sowing (Petrović et al., 2008), level of water (Pawlonka &

Skrzyczyńska, 2004), fertilizer pesticide application (Füleky, 2008). In wheat and other cereal species yield losses under average weediness can reach about 30% (Mason et al., 2007). One of the most important elements of the crops protection is weed control. Time of sowing and density (Walter et al., 2002) of wheat is important for tillering, enough to suppress weed in spring season and reduce yield losses caused by late sowing, low temperature during winter and early spring time. Wheat density can influence the reduction of weed dry matter more than 60% (Lemerle et al., 1996) and reduction to weed structure (Korres & Froud-Williams, 2002). The application of herbicides do not influence the wheat density and wheat yield (Kristensen et al., 2008). Adequate wheat growing contributes to competitiveness with weed. Crop shading ability contributes to weed control by decreasing chemical application and environmental protection (Lemerle et al., 1996). Increase in wheat density and uniformity can help in suppression of weeds (Olsen et al., 2006). Wheat cultivars differ in competitiveness against weeds (Avramovic & Knežević, 2007) therefore the selection of cultivar is efficient preventive method of weed control (Hansen et al., 2008) and expression of economic traits. Wheat grain yield, as well as yield of other small grains, is in dependence of interaction of genotypes and environment (Oettler et al., 2006; Pepo, 2007) mineral elements availability (Kovačević et al., 2005) agro-ecological condition (Dreznar et al., 2007) precipitation (Balogh et al., 2006) crop weediness (Fodor & Pálmai, 2008; Knežević et al., 2008). Also optimal application of fertilizer will increase vigour and competitiveness of cereal crop (Paunović et al., 2007). Grain yield and yield components and protein content increased with increasing NP fertilizer rates (Zečević et al., 2012). The increasing of nitrogen rate resulted in increasing of biomass formation both of cereal plants and weeds (Jolánkai et al., 2006; Knežević et al., 2007).

The aim of this paper is investigation of the effect of different rate of nitrogen fertilization to weed communities of winter winter wheat cultivars and variation of grain yield.

## Materials and methods

The Lazarica and Julija Mono wheat cultivars grown under different regime of nitrogen nutrition were studied for weediness and grain yield. Cultivars were grown on long term experimental field of Center for Small Grains in Kragujevac on control variant (unfertilized), second variant with nitrogen application (two sub variants with different rate of N)  $2N_1$  ( $80 \text{ kg P ha}^{-1}$ ) and  $2N_2$  ( $120 \text{ kg N ha}^{-1}$ ), third variant with NP fertilizer with two different rate of N application:  $3N_1$  ( $80-100 \text{ kg NP ha}^{-1}$ ) and  $3N_2$  ( $120-100 \text{ kg NP ha}^{-1}$ ) and fourth variant  $4N_1$  ( $80-60 \text{ kg NP ha}^{-1}$ ) and  $4N_2$  ( $120-60 \text{ kg NP ha}^{-1}$ ). The analysis weed community and grain yield of wheat cultivars were conducted on four basic variants of soil fertilization and three replications. Unfertilized plots belonged to the smonitza type of soil in the process of degradation with pH of 6.03 to 6.10 in water and 4.76 to 4.84 in KC1. Floristic structure was estimated by method Braun-Blanquet (1964).

## Results and discussion

The weed species in wheat cultivated under different regime of fertilization were identified. Different rates of nitrogen and phosphorus were influenced differences of weediness (tab. 1. and tab. 2). The lowest weediness registered on the plots of wheat crops where applied the highest rate of nitrogen. In each variant of applied fertilizer rates were registered between 7 and 11 weed species in both tested wheat cultivars Lazarica and Julija Mono. Mainly in all variants of nitrogen fertilization the weed species *Cirsium arvense* L., *Consolida regalis* Gray, *Convolvulus arvensis* L., *Polygonum aviculare* L. was dominant (tab.1 and tab 2).

The study of floristic structure determined that weeds community represents a fragment of *Consolida regalis-Polygonum aviculare* association having in it's composition 13 species. The weed species *Consolida regalis* Gray, *Convolvulus arvensis* L., *Polygonum aviculare* L., *Cirsium arvensis* L., *Polygonum convolvulus* L. were dominant on fertilized plots. These results are in agreement with previous results presented by Knezevic et al. (2008).

Different rate of nitrogen fertilization had positive effect on wheat crop density and no influence to the weeding level and botanical composition of the community. On control plots it was observed low density of crops. Fertilized plots characterized by presence of similar floristic composition of weed species in analyzed wheat cultivars, and registered less weed species as well as their different density. The increase of fertilizers rate had influence to the increase of the wheat plants tillering, the more development of the wheat plants and higher yields. Also, in these conditions the density of the weeds was reduced.

Other authors registered the most prevalent weed species *Capsella bursa-pastoris* L. and *Taraxacum officinale* F.H. Wigg (Harker et al. 2000). Density of weeds is dependent on weed species and soil properties. So, negative cross-correlation between the density of *Viola arvensis* Murray and clay content, while, density of *Lamium purpureum* L. was positively cross-correlated with the phosphorus content in the soil (Walter et al., 2002).

The interaction between cultivar and N supply had significant impact on yield. In this investigation we analyzed influence of weeds and fertilizer to variation of grain yield in wheat. The highest grain yield was found for wheat cultivars on the plots where the highest rates of fertilizer were applied. Cultivar Julija Mono expressed the highest grain yield ( $4990 \text{ kg ha}^{-1}$ ) and low weediness in fertilizers variant of  $3N_2$  ( $120 \text{ kg N ha}^{-1} + 100\text{kg P ha}^{-1}$ ) what could be advantage for cultivation (tab. 3). On the same variant of fertilization  $3N_2$  ( $120 \text{ kg N ha}^{-1} + 100\text{kg P ha}^{-1}$ ) other investigated wheat cultivar Lazarica achieved the highest yield ( $3724 \text{ kg ha}^{-1}$ ). Grain per square meter were positively correlated with grain yield and influenced by kernels per spikelet, a measure of fertility (Bennett et al., 2012).

Tab. 1. Weed species in wheat in control variant (no fertilized plots) 1N<sub>1</sub>, and in a variant with application of different NP fertilizer rates 2N<sub>1</sub> (80kg N ha<sup>-1</sup>), 3N<sub>1</sub> (80-100kg NP ha<sup>-1</sup>) 4N<sub>1</sub> (80-100kg NP ha<sup>-1</sup>)  
 Врсте корова у пшеници у контролној варијанти (без додатних паруена) 1N<sub>1</sub>, и на варијантама са применом различитих доза азотног NP додатка 2N<sub>1</sub> (80kg N ha<sup>-1</sup>), 3N<sub>1</sub> (80-100kg NP ha<sup>-1</sup>) 4N<sub>1</sub> (80-100kg NP ha<sup>-1</sup>)

	1N <sub>1</sub> (control)	2N <sub>1</sub> (80kg N ha <sup>-1</sup> )	3N <sub>1</sub> (80-100kg NP ha <sup>-1</sup> )	4N <sub>1</sub> (80-100kg NP ha <sup>-1</sup> )
	Lazarica	Julija Mono	Lazarica	Julija Mono
<i>Agropyrum repens</i> L.	-	1.1	-	+1
<i>Consolida regalis</i> Gray	+1	1.1	+1	1.1
<i>Convolvulus arvensis</i> L.	-	+1	-	+1
<i>Linaria vulgaris</i> Mill.	+1	-	1.1	-
<i>Sorghum halepense</i> L.	-	-	-	-
<i>Viola arvensis</i> Murray	r	-	r	+1
<i>Polygonum aviculare</i> L.	+1	1.1	r	+1
<i>Polygonum convolvulus</i> L.	-	+1	-	+1
<i>Chenopodium album</i> L.	-	-	-	-
<i>Veronica opaca</i> Fries	r	-	-	-
<i>Cirsium arvense</i> L.	+1	1.1	+1	+1
<i>Rubus caesius</i> L.	1.1	-	-	-
<i>Mercurialis</i>	r	-	-	-
Total no. of species:	11	9	8	9
			7	9
			9	9
			10	10

Tab. 2. Weed species in wheat on controls variant (no fertilized plots) 1N<sub>2</sub>, and on variant with application of different NP fertilizer rates 2N<sub>2</sub> (120 kg N ha<sup>-1</sup>) 3N<sub>2</sub> (120-100 kg NP ha<sup>-1</sup>) 4N<sub>2</sub> (120-60 kg NP ha<sup>-1</sup>)  
*Врсме корова у пшеници ю контролној варијанти (без хидратних натријена) 1N<sub>2</sub>, и на варијанту са приметом различитих доза азотног NP хидратива 2N<sub>2</sub> (80kg N ha-1), 3N<sub>2</sub> (80-100kg NP ha-1) 4N<sub>2</sub> (80-100kg NP ha-1)*

	1N <sub>2</sub> (control)			2N <sub>2</sub> (120kg N ha <sup>-1</sup> )			3N <sub>2</sub> (120-100kg NP ha <sup>-1</sup> )			4N <sub>2</sub> (120-60 kg NP ha <sup>-1</sup> )			
	Lazarica	Julija Mono	Lazarica	Julija Mono	Lazarica	Julija Mono	Lazarica	Julija Mono	Lazarica	Julija Mono	Lazarica	Julija Mono	
<i>Agropyrum repens</i> L.	+.1	1.1	+.1	1.1	r	+.1	-	+.1	-	r	1.1	+.1	1.1
<i>Consolida regalis</i> Gray	+.1	-	-	+.1	1.1	-	1.1	+.1	-	+.1	1.1	+.1	1.1
<i>Convolvulus arvensis</i> L.	+.1	-	+.1	1.1	-	+.1	1.1	-	+.1	+.1	1.1	+.1	1.1
<i>Linaria vulgaris</i> Mill.	-	-	+.1	1.1	r	-	-	-	-	-	-	-	-
<i>Sorghum halepense</i> L.	-	-	r	-	r	-	r	-.1	+.1	-.1	+.1	-.1	+.1
<i>Viola arvensis</i> Murray	-	-	r	+.1	-	r	-.1	+.1	r	+.1	1.1	+.1	1.1
<i>Polygonum aviculare</i> L.	+.1	-	+.1	1.1	r	1.1	+.1	-	+.1	+.1	1.1	+.1	1.1
<i>Polygonum convolvulus</i> L.	-	+.1	+.1	-	+.1	-	+.1	-.1	+.1	+.1	1.1	+.1	1.1
<i>Chenopodium album</i> L.	-	-	-	-	-	-	-	-	-	r	-	-	-
<i>Veronica opaca</i> Fries	r	-	-	r	-	-	-	-	-	-	-	-	-
<i>Cirsium arvense</i> L.	+.1	1.1	+.1	1.1	-	+.1	1.1	+.1	1.1	+.1	1.2	r	1.1
<i>Rubus caesius</i> L.	-	-	+.1	-	-	-	+.1	1.1	+.1	1.1	-	r	-
<i>Mercurialis</i>	-	-	-	r	-	r	-	-	-	-	-	-	-
Total no. of species:	7	7	11	10	8	9	10	9	10	9	10	9	10

The least grain yield was found on control variant for both of analyzed wheat cultivars Lazarica ( $1880 \text{ kg ha}^{-1}$ ) and Julija Mono ( $1740 \text{ kg ha}^{-1}$ ) tab. 3. Cultivar Julija Mono had higher grain yield ( $2586 \text{ kg ha}^{-1}$ ) on plot fertilized by  $80 \text{ kg N ha}^{-1}$  (variant  $2N_1$ ) than cultivar Lazarica ( $2420 \text{ kg ha}^{-1}$ ). Also on variant  $2N_2$  ( $120 \text{ kg N ha}^{-1}$ ) cultivar Julija Mono had higher grain yield ( $2680 \text{ kg ha}^{-1}$ ) than cultivar Lazarica ( $2470 \text{ kg ha}^{-1}$ ).

Tab. 3. Average yield of grain ( $\text{kg ha}^{-1}$ ) of examined wheat cultivars under the different variants of fertilization

*Просјечан принос зрна ( $\text{kg ha}^{-1}$ ) испитиваних пшеничних сорти под различитим варијантама ђубрења*

Cultivar / Сорта	Lazarica		Julija Mono	
	Plot Парцела	Hectare	Plot Парцела	Hectare
Variants / Варијанте				
Control 1	940	1880	890	1780
Control 2	955	1910	870	1740
$2N_1$ ( $80 \text{ kg N ha}^{-1}$ )	1210	2420	1293	2586
$2N_2$ ( $120 \text{ kg N ha}^{-1}$ )	1235	2470	1340	2680
$3N_1$ ( $80 \text{ kg N ha}^{-1} + 100 \text{ kg P ha}^{-1}$ )	1796	3592	2192	4384
$3N_2$ ( $120 \text{ kg N ha}^{-1} + 100 \text{ kg P ha}^{-1}$ )	1862	3724	2495	4990
$4N_1$ ( $80 \text{ kg N ha}^{-1} + 60 \text{ kg P ha}^{-1}$ )	1565	3130	1996	3992
$4N_2$ ( $120 \text{ kg N ha}^{-1} + 60 \text{ kg P ha}^{-1}$ )	1645	3290	2178	4356

In general, increasing rates of NPK fertilization caused only a significant increase in the yields. Similar results obtained in the crop rotation at lower rates  $150\text{--}300 \text{ kg NPK ha}^{-1}$  (Füleky, 2008). Increasing assimilate allocation to the reproductive primordia indirectly increase total crop photosynthesis and intensive development growth of vegetative reproductive organs in wheat. However, total photosynthesis has increased as a result of an increase in leaf area, daily duration of photosynthesis or leaf area duration (Richards, 2000). Even small increases in the rate of net photosynthesis can translate into large increases in biomass and hence yield, since carbon assimilation is integrated over the entire growing season and crop canopy.

## Conclusion

The results of investigation showed differences of weediness in wheat cultivars and specific interaction genotype/nitrogen rate. The increasing in fertilizer rate affected grain yield as well abundance of weed species. The higher rates of nitrogen fertilizer rate have effect to efficient competition of wheat plants and composition of weed species which leads to increase of grain yield. The highest grain yield in wheat Julija Mono ( $4990 \text{ kg ha}^{-1}$ ) was found in variant  $3N_2$  ( $120 \text{ kg N ha}^{-1} + 100 \text{ kg P ha}^{-1}$ ). Also tested cultivar Lazarica achieved the highest yield ( $3724 \text{ kg ha}^{-1}$ ). Breeding of wheat cultivars for improving grain yield require complex work in direction to

improving several types of abiotic stress as well tolerance to drought, high temperatures, high irradiance, and nutrient toxicities or deficiencies. A strategy is then described where a specific environment is targeted and appropriate germplasm adapted to the chosen environment, based on extensive definition of the morpho-physiological and molecular mechanisms of tolerance of the parents. In wheat grown under different application of nitrogen fertilizer identified different weed species while prevail weed species were *Cirsium arvense* L., *Consolida regalis* Gray, *Convolvulus arvensis* L., *Polygonum aviculare* L. For weed management strategy in wheat should be optimize technological growing measures. One of the strategies to reduce weediness based on increasing of sowing density in conventional agriculture and it is a way to reduce herbicide application levels and further this strategy may have other positive influence to environment including fuel consumption and carbon dioxide (CO<sub>2</sub>) production.

### Acknowledgements

Authors gratefully acknowledge the financial support by the Ministry of Education, Science and Technological development of Republic Serbia, Belgrade, Project Code TR-31092.

### References

- Avramovic, S. & Knezevic, D. (2007). Weed community in wheat and triticale crops. *Proceedings of Int. Conference of Agricultural faculties, Cacak*, 2, 129-139.
- Balogh, A., Pepo, P. & Hornok, M. (2006). Interactions of crop year, fertilization and variety in winter wheat management. *Cereal Research Communications*, 34(1), 389-392.
- Bennett, D., Izanloo, A., Reynolds, M., Kuchl, H., Langridge, P. & Schnurbusch, T. (2012). Genetic dissection of grain yield and physical grain quality in bread wheat (*Triticum aestivum* L.) under water-limited environments. *Theor. Appl. Genet.*, 125(2), 255-271.
- Braun-Blanquet, J. (1964). *Pflanzensoziologie*. Wien-New York.
- Drezner, G., Dvojkovic, K., Horvat, D., Novoselovic, D. & Lalic, A. (2006). Environmental impacts on wheat agronomic and quality traits. *Cereal Research Communications*, 35(2), 357-360.
- Fodor, L. & Pálmai, O. (2008). The influence of nitrogen fertilization and sowing time on the weediness of winter wheat. *Cereal Research Communications*, 36, 1159-1162.
- Füleky, G. (2008). Results of a 30-year-old fertilisation experiment. *Acta Agronomica Hungarica*, 56, 265-273.
- Harker, K.N., Baron, S. V., Chanasyk, S. D., Naeth, M. A. & Stevenson, F. C. (2000). Grazing intensity effects on weed populations in annual and perennial pasture systems. *Weed Science*, 48(2), 231-238.
- Jolánkai, M., Szentpétery, Zs. & Hegedűs, Z. (2006). Pesticide Residue discharge dynamics in wheat grain. *Cereal Research Communications*, 34, 505-509
- Knežević, D., Paunović, A., Madić, M. & Djuki, N. (2007). Genetic analysis of nitrogen accumulation in four wheat cultivars and their hybrids. *Cereal Research Communications*, 35, 633-636.

- Knežević, D., Avramović, S. & Branković, G. (2008). Weed community in wheat crops and influence to grain yield under different regime of nutrition. *Acta Herbologica*, 17, 95-106.
- Kristensen, L., Olsen, J. & Weiner, J. (2008). Crop density sowing pattern, and nitrogen fertilization effect on weed suppression and yield in spring wheat. *Weed Science*, 56(1), 97-102.
- Korres, N.E. & Froud-Williams, R.I. (2002). Effects of winter wheat cultivars and seed rate on the biological characteristics of naturally occurring weed flora. *Weed Research*, 42(6), 417-428.
- Kovačević, V., Petosić, D. & Josipović, M. (2005). Potassium availability in hydromorphic soils of Eastern Croatia. *Cereal Research Communications*, 33(1), 247-250.
- Lemerle, D., Verbeek, B., Cousens, R. D. & Coombes, N.E. (1996). The potential for selecting wheat varieties strongly competitive against weeds. *Weed Research*, 36(6), 505-513.
- Mason, H.E., Navabi, A., Frick, B.L., O'Donovan, J. T. & Spaner, D.M. (2007). The Weed Competitive Ability of Canada Western Red Spring Wheat Cultivars Grown under Organic Management. *Crop.Sci.*, 47(3), 1167-1176
- Márton, L. (2008). Impact of rainfall, liming, nitrogen (N), phosphorus (P<sub>2</sub>O<sub>5</sub>), potassium (K<sub>2</sub>O), calcium (CaO), magnesium (MgO) mineral fertilization on triticale ( $\times$  Triticosecale Wittmack) yield in a monoculture in Hungary. *Cereal Research Communications*, 36(2), 333-341
- Olsen, J., Kristensen, L. & Weiner, L. (2005). Effects of density and spatial pattern of winter wheat on suppression of different weed species. *Weed Science*, 53(5), 690-694.
- Oettler, G., Wiethölter, S. & Horst, W.J. (2000). Genetic parameters for agronomic traits of triticale and other small-grain cereals grown on aluminium-toxic soil in southern Brazil. *Plant breeding*, 119(3), 227-231.
- Paunović, A., Madić, Milomirka, Knežević, D. & Bokan, N. (2007). Sowing density and nitrogen fertilization influences on yield components of barley. *Cereal Res. Commun.*, 35(2), 901-904.
- Pawlak, Z. & Skrzyczyńska, J. (2004). The effect of selected agrotechnical factors and weather conditions on the weeding of winter triticale. *Annales UMCS, Sec. E*, 59(3), 1037-1047.
- Pepo, P. (2007). The role of fertilization and genotype in sustainable winter wheat (*Triticum aestivum* L.) production. *Cereal Res. Comm.* 35(2), 917-920.
- Petrović, S., Marić, S., Guberac, V., Drezner, G. & Eded A. (2008). Influence of environmental conditions and sowing rates on winter wheat yield. *Cer.Res.Comm.*, 36(2), 307-1310.
- Richards, R. A. (2000). Selectable traits to increase crop photosynthesis and yield of grain crops. *J. Exp. Bot.*, 51(suppl. 1), 447-458.
- Sepp, K., Kanger, J. & Särekanno, M. (2009). Influence of soil tillage methods on the weediness and yields of spring wheat, spring barley and field pea in organic crop rotation. *Agronomy Research*, 7(Special issue I), 477-484.
- Walter, A M., Christensen, S. & Simmelsgaard, S. E. (2002). Spatial correlation between weed species densities and soil properties. *Weed Research*, 42(1), 26-38.

Zečević, V., Bošković, J., Knežević, D. & Mićanović, D. (2012). The effect of fertilization and liming on some grain quality properties of wheat. *Review on Agriculture and Rural Development. "Science for Rural Areas" XI. Wellmann International Scientific Conference, Hódmezővásárhely (Hungary), 10th May, 2012. Conference proceedings*, pp. 470-475.

## Принос пшенице и присуство корова при примени различитих доза азотног ћубрива

Десимир Кнежевић<sup>1</sup>, Димитрије Марковић<sup>2</sup>,  
Сретенка Марковић<sup>2</sup>, Данијела Кондић<sup>2</sup>

<sup>1</sup> Универзитет у Приштини, Пољопривредни факултет, Косовска Митровица-Лешак, Србија

<sup>2</sup>Пољопривредни факултет, Универзитет у Бањој Луци, Босна и Херцеговина

### Сажетак

У овом раду је представљено изучавање приноса зрна и појава коровских врста у усевима две сорте озиме пшенице (Лазарица и Јулија Моно), гајених у условима различитог режима минералне исхране. Изучавања су обављена на дугогодишњем огледу са четири варијанте ћубрења са по две варијанте са различитом дозом примене азотног ћубрења: контролна варијанта (без ћубрења), друга варијанта примене азота са две подваријанте са различитом дозом азота  $2N_1$  ( $80 \text{ kg N ha}^{-1}$ ) и  $2N_2$  ( $120 \text{ kg N ha}^{-1}$ ), трећа варијанта са две подваријанте  $3N_1$  ( $80 \text{ kg N ha}^{-1} + 100 \text{ kg P ha}^{-1}$ ) и  $3N_2$  ( $120 \text{ kg N ha}^{-1} + 100 \text{ kg P ha}^{-1}$ ) и четврта варијанта са две подваријанте  $4N_1$  ( $80 \text{ kg N ha}^{-1} + 60 \text{ kg P ha}^{-1}$ ) and  $4N_2$  ( $120 \text{ kg N ha}^{-1} + 60 \text{ kg P ha}^{-1}$ ). Оцена је урађена у три понављања на свим варијантама ћубрења. Метода Braun-Blanquet (1964) је коришћена за оцену флористичког састава у усевима две сорте пшенице. Регистровано је 13 различитих коровских врста са различитом бројношћу, која је показала одређену зависност са примењеним дозама азотног ћубрива. На нећубреним парцелама је установљено веће присуство корова. Повећавање дозе азотног ћубрива на парцелама је утицало на већу густину усева пшенице и смањен интензитет коровских врста. Коровске врсте *Consolida regalis* Gray, *Convolvulus arvensis* L., *Polygonum aviculare* L., *Cirsium arvense* L., *Agropyrum repens* L. и *Polygonum convolvulus* L. су биле доминантне на ћубреним парцелицама. Изучаван је принос зрна пшенице и установљено је варирање зависно од исхране. На парцелама без додавања азота принос зрна је био најнижи код сорте Лазарица ( $1880 \text{ kg ha}^{-1}$ ), и Јулија Моно ( $1740 \text{ kg ha}^{-1}$ ). Највећи приноси зрна су добијени на парцелама са варијантом

ћубрења  $3N_2$  ( $120 \text{ kg N ha}^{-1} + 100\text{kg P ha}^{-1}$ ) и то код Лазарице ( $3724\text{kg ha}^{-1}$ ) и Јулије Моно ( $4990\text{kg ha}^{-1}$ ) а на овој варијанти је установљена и најмања закоровљеност. Ово указује да оптимизација примене ћубрења доприноси повећању густине усева пшенице, већој компетитивности са коровским врстама, што утиче на мање присуство корова и смањену потребу употребе хербицида, чиме се значајно доприноси смањењу трошкова енергије и очувању животне средине.

*Кључне речи:* компетитивност, закоровљеност, пшеница, спољашња средина, принос зрна, доза ћубрења

Desimir Knežević

E-mail address:

[desimir.knezevic@pr.ac.rs](mailto:desimir.knezevic@pr.ac.rs)